

Changes in the central ANF-system of renovascular hypertensive rats

UDO BAHNER, HELMUT GEIGER, MIKLÓS PALKOVITS, DETLEV GANTEN, BURKHARD KLOTZ, and AUGUST HEIDLAND

Medical Clinic of the University, Division of Nephrology, Würzburg, Federal Republic of Germany; First Department of Anatomy, Semmelweis University, Medical School, Budapest, Hungary; and Department of Pharmacology, University of Heidelberg, Heidelberg, Federal Republic of Germany

Changes in the central ANF-system of renovascular hypertensive rats. The central atrial natriuretic peptides (ANF)-system was investigated in volume-dependent one-kidney, one-clip (1K1C) and renin-dependent two-kidney, one-clip (2K1C) renovascular hypertensive rats by radio-immunological measurement of ANF concentration in 18 selected brain areas. Significant changes were found in nine brain areas of 1K1C and in eight brain areas of 2K1C hypertensive rats. Except unidirectional changes in the organum vasculosum laminae terminalis and the supraoptic nucleus, ANF concentration was changed in the opposite direction in all other brain areas, with an activation of the central ANF system in 1K1C and an inhibition in 2K1C hypertension. The localization of the alterations (circumventricular organs, anteroventral third ventricle region, hypothalamo hypophyseal system, brain stem) implies major differences in the central regulation of blood pressure and electrolyte and fluid homeostasis between these two models. The activation of the central ANF system in 1K1C hypertension may be a compensatory mechanism to prevent further increments in blood pressure and plasma volume. In contrast, the depression of the central ANF system in 2K1C hypertension may promote the elevation of the blood pressure.

Since Goldblatt et al [1] demonstrated the development of arterial hypertension following renal ischemia, the pathogenesis of renovascular hypertension has been intensively investigated. However, the mechanism of the hypertension in renal artery stenosis is still not completely understood. Clipping of one renal artery regularly induces hypertension, but there are differences between the two-kidney, one-clip (2K1C) model, in which the contralateral kidney is untouched, and the one-kidney, one-clip (1K1C) model, in which the contralateral kidney is removed. The 2K1C model seems to be dependent on the renin-angiotensin system, while the 1K1C model is suggested to be volume expanded [2–5].

There is an increasing number of indications that the atrial natriuretic factor (ANF), a peptide hormone that is intimately involved in the regulation of renal and cardiovascular homeostasis [6–8], may play an important role in the pathomechanism

of renovascular hypertension. In 1K1C rats plasma ANF levels are positively correlated with blood pressure and negatively correlated with ANF concentration in the left atrium [9]. Additional infusion of synthetic ANF results in a reduction of blood pressure and increased diuresis and natriuresis in 1K1C rats [10]. Unclipping of the remaining kidney is accompanied by a reversal of hypertension, which might be partially due to the still higher plasma ANF levels [9]. Based on these findings, it was suggested that the increased plasma ANF level in 1K1C hypertension reflects a compensatory mechanism preventing further increments in blood pressure and plasma volume. In 2K1C rats plasma ANF level is unchanged, but chronic administration of synthetic ANF also decreases blood pressure in renin-dependent 2K1C hypertension [11]. The hypotensive effect of ANF in this model may be related to the inhibition of renin release and the consequent decrease of plasma angiotensin II and aldosterone levels [12].

However, the mechanisms underlying the shifting pattern of hemodynamic and humoral events in the various stages and different forms of renal hypertension may not only be localized in the periphery but may be combined with regulatory mechanisms in the central nervous system. Thus, it was shown that 1K1C hypertension is associated with an increased activity of the sympathetic nervous system [13,14], and that ablation of the anteroventral third ventricle (AV3V) region prevents the development of 2K1C hypertension [15,16]. Since the central ANF system is known to be important for the regulation of electrolyte and fluid homeostasis and blood pressure [17–21], it may also be involved in the pathophysiology of renovascular hypertension. To elucidate this question, we have investigated the central ANF system in the early maintenance phase of 1K1C and 2K1C renovascular hypertension, when differences in the pathological disorders could be expected. For this purpose, the concentration of ANF was measured radioimmunologically in 18 selected brain areas of 1K1C and 2K1C hypertensive rats and compared with the respective controls. Previous studies have shown that the concentration of ANF in specific brain areas is a sensitive parameter for alterations in the electrolyte and fluid homeostasis and blood pressure [22–24].

Received for publication March 30, 1990
and in revised form August 3, 1990
Accepted for publication August 3, 1990

© 1991 by the International Society of Nephrology

Methods

Protocol

Renovascular hypertension was produced in male Sprague-Dawley rats (Charles River Wega GmbH, Sulzfeld, FRG; weight 80 to 100 g; age 35 to 40 days) under sodium pentobarbital anesthesia (60 mg/kg body wt i.p.) by partial constriction of the left renal artery with a silver clamp of 0.20 mm internal diameter. In the 1K1C hypertension group the contralateral kidney was removed simultaneously, whereas it was left untouched in the 2K1C hypertension group. Animals subjected to sham operation, in which a silver clamp was temporarily placed around the left renal artery and then taken away and the right kidney was removed (1K1C control group) or left untouched (2K1C control group), served as normotensive controls. The animals were housed at a constant room temperature of $22^{\circ} \pm 1^{\circ}$ C and a humidity of $60\% \pm 4\%$, with illumination from 6.00 a.m. to 6.00 p.m. Systolic blood pressure was measured indirectly once a week by means of the tail cuff technique. The criterion for classifying renovascular hypertensive and sham operated animals were as follows: clipped animals having a systolic blood pressure higher than or equal to 160 mm Hg (about 70% of all clipped animals) were classified as renovascular hypertensive rats; all animals selected for the 2K1C hypertension group had to have a four fold increased plasma renin activity; sham operated animals having a systolic blood pressure less than or equal to 120 mm Hg (about 90% of all sham operated animals) were classified as normotensive.

Determination of blood parameters

The blood parameters were determined separately in 10 animals per experimental group. The animals were chosen randomly and moved to a quiet room. One at a time they were transferred to an adjacent room.

For determination of hematocrit, plasma osmolality, serum sodium, potassium and chloride concentration blood was collected by puncture of the aorta under sodium pentobarbital anesthesia (60 mg/kg body wt i.p.). Hematocrit was determined by an automatic analyzer (ELT 8, Fa. Moltronic, Heidelberg, FRG), plasma osmolality by freezing point depression (Osmomat, Fa. Gonotec, Berlin, FRG), serum sodium and potassium by flame photometry (AFM 2, Eppendorf Co., Hamburg, FRG) and chloride by mercurimetry (Chlorimeter, Eppendorf Co., Hamburg, FRG).

For determination of plasma renin activity (PRA) and angiotensin II (Ang II) concentration rats were rapidly anesthetized with ether, a laparotomy performed, and blood collected from the abdominal aorta. The interval between the beginning of the exposure to ether and the end of the blood withdrawal never exceeded 150 seconds. Blood was collected into prechilled (4° C) polypropylene tubes, which contained 5% of a solution of 125 mM EDTA for determination of PRA or 5% of a solution of 125 mM EDTA and 26 mM orthophenanthroline for determination of Ang II concentration. After centrifugation (6000 g, 10 min, at 4° C) plasma was stored at -20° C until assay. PRA was determined by radioimmunological measurement of generated angiotensin I [25] and Ang II was measured by direct radioimmunoassay in unextracted plasma [26].

Plasma aldosterone concentration was measured in heparinized, unextracted plasma by a commercial aldosterone radioimmunoassay (Biermann Diagnostica, Bad Nauheim, FRG).

Plasma arginine vasopressin concentration was measured after extraction from plasma by radioimmunoassay as described elsewhere [27].

For determination of plasma norepinephrine concentration blood was collected during the first 3 seconds after rapid decapitation into prechilled (4° C) polypropylene tubes containing EDTA. The measurement was performed by high-pressure liquid chromatography using a Beckman 114 M pump and a Chromatopak C-18 column. The samples were eluted in an isocratic system with a 0.06 M phosphate buffer at a flow rate of 0.5 ml/min. The detection was carried out electrochemically at + 650 mV towards an Ag/AgCl electrode [28].

For determination of plasma ANF concentration, blood was taken from the aorta, collected in polypropylene tubes containing 500 KIU aprotinin (Trasylol[®], Bayer AG, Leverkusen, FRG), and immediately centrifuged (2.000 g, 10 min, at 4° C). The plasma was stored at -70° C until assay. ANF was extracted from plasma according to the method of Larose et al [29] modified in some details. Plasma sample (1 ml) acidified to pH 3 to 4 with 2 N HCL was applied to a Sep Pak C18 reverse-phase octadecyl silane cartridge column (Water Associates, Milford, Massachusetts, USA) that had been activated previously with 100% methanol (10 ml) followed by 5% acetic acid (10 ml). After washing with trifluoroacetic acid (TFA, 3×5 ml), the peptide was eluted with a mixture of 60% acetonitrile in 0.1% TFA (2 ml). The eluates were evaporated to dryness with nitrogen, and the residue was reconstituted with 1 ml of RIA buffer (50 mM potassium phosphate buffer, pH 7.4, containing 0.1% BSA, 0.02 Na₂N₃, 50 mM NaCl, 0.1% Triton X-100, 50 μ M phenylmethylsulfonyl fluoride and 10 μ M aprotinin) for ANF determination by radioimmunoassay.

Tissue dissection and extraction of brain ANF

The animals were decapitated between 8.00 and 10.00 a.m., brains were quickly removed and frozen on dry ice. Serial coronal sections with 300 μ m thickness were cut in a cryostat on -10° C and 18 brain regions (Table 2) were removed by the micropunch technique [30]. The brain regions represented those functional systems which were mainly involved in the regulation of electrolyte and fluid homeostasis and blood pressure. Tissue pellets of identical brain nuclei of four animals were pooled, and six assay sets per group were formed (4×24 rats). The samples were collected in 1.5 ml conical Eppendorf tubes containing 60 μ l 0.1 N HCL and put in a boiling water bath for 10 minutes. After addition of 100 μ l RIA-buffer the samples were homogenized and 20 μ l aliquots of the homogenates were removed for protein determination according to Lowry et al [31]. The remaining sample was centrifuged at 2000 g for 15 minutes at 4° C and 100 μ l of the supernatant fluid were used for ANF determination by radioimmunoassay.

Measurement of plasma ANF and brain ANF concentration

For determination of plasma ANF and brain ANF concentration a commercial radioimmunoassay kit was used (Peninsula Labs., INC, Belmont, California, USA; RIK 8798). The rabbit antihuman ANF (99-126) antiserum of this kit cross reacts with the different atrial natriuretic peptides as follows: 100% with

Table 1. Characteristics of 1K1C and 2K1C renovascular hypertensive rats and their control rats (mean \pm SEM)

	1K1C control	1K1C	2K1C control	2K1C
No.	10	10	10	10
Body weight g	262.0 \pm 8.5	246.5 \pm 9.5	266.0 \pm 8.5	232.5 \pm 7.0
Systolic blood pressure mm Hg	112.5 \pm 3.5	178.0 \pm 4.5 ^c	116.5 \pm 3.5	175.0 \pm 5.0 ^c
Water intake % of body wt	15.5 \pm 1.5	10.8 \pm 2.6 ^a	15.0 \pm 0.5	23.0 \pm 1.2 ^a
Urine volume % of body wt	7.3 \pm 0.9	5.3 \pm 0.6	8.0 \pm 0.4	11.0 \pm 1.3 ^a
Hematocrit %	41.7 \pm 0.5	38.1 \pm 0.4 ^a	42.1 \pm 0.4	46.1 \pm 0.4 ^a
Plasma sodium concentration mmol/liter	147.7 \pm 1.0	141.2 \pm 1.6 ^b	150.8 \pm 0.6	148.1 \pm 1.4
Plasma potassium conc. mmol/liter	4.4 \pm 0.1	4.5 \pm 0.2	4.5 \pm 0.2	4.4 \pm 0.2
Plasma chloride conc. mmol/liter	99.5 \pm 1.3	98.0 \pm 0.4	103.7 \pm 0.7	101.7 \pm 1.5
Plasma osmolality mOsm/kg water	315.4 \pm 1.4	302.4 \pm 2.0 ^b	318.6 \pm 2.4	316.2 \pm 1.9
Plasma renin activity ng Ang I/ml/hr	1.5 \pm 0.2	1.4 \pm 0.2	1.9 \pm 0.4	9.8 \pm 0.6 ^c
Plasma angiotensin II conc. fmol/ml	152.3 \pm 23.1	147.1 \pm 16.2	150.5 \pm 21.0	956.5 \pm 93.5 ^c
Plasma aldosterone conc. fmol/ml	81.0 \pm 19.7	67.2 \pm 16.2	94.5 \pm 20.2	486.0 \pm 81.0 ^b
Plasma norepinephrine conc. pmol/liter	1.0 \pm 0.3	3.5 \pm 0.4 ^b	1.3 \pm 0.2	1.4 \pm 0.3
Plasma arginine vasopressin conc. pg/ml	1.0 \pm 0.1	1.1 \pm 0.2	1.1 \pm 0.1	2.6 \pm 0.4 ^b
Plasma ANF concentration pg/ml	7.2 \pm 0.8	29.6 \pm 2.5 ^c	8.0 \pm 1.3	12.6 \pm 1.8

^a $P < 0.05$, ^b $P < 0.01$, ^c $P < 0.001$, significant difference from the renovascular hypertensive rats to their respective control rats

human ANF (99-126), rat ANF (99-126) and rat ANF (102-126), 90% with rat ANF (101-126), 88% with rat ANF (103-126), 57% with ANF (108-126), 50% with rat ANF (111-126), 27% with rat ANF (103-125), 10% with rat ANF (102-125) and 3% with rat ANF (103-123). The ANF antiserum does not cross react with somatostatin, oxytocin, arginine⁸-vasopressin and brain natriuretic polypeptide porcine (BNP). In conclusion, the antiserum recognizes nearly completely the active forms of ANF as well in the plasma (rat ANF (99-126)) [7,32] as in the brain (rat ANF (102-126) and rat ANF (103-126)) [33,34].

The reconstituted plasma extracts (100 μ l), respectively, the brain extracts (100 μ l) were incubated with the same volume of the antiserum. Synthetic human ANF (99-126) ranging from 1 to 128 pg/tube was incubated as standard. After incubation for 24 hours at 4°C, 100 μ l of the ANF tracer ¹²⁵I-human ANF (99-126) was added. After again 24 hours incubation at 4°C, goat anti-rabbit IgG serum and normal rabbit serum (100 μ l, dissolved in RIA buffer) were added and the mixture was incubated at room temperature for two hours. Then 500 μ l RIA buffer was added, the contents were vortex mixed and then centrifuged (2000 g, 20 min, at 20°C). After centrifugation the supernatants were aspirated and the precipitants were counted in a gamma counter. The sensitivity of the assay was 1.5 pg/tube. The recovery of added ANF standard was 78.7 \pm 3.3% (mean \pm SEM, $N = 6$) and the intraassay and interassay variations were 9% and 14%, respectively. Serial dilutions of plasma and brain extracts showed parallelism to the synthetic ANF standards.

Statistical analysis

The results are expressed as mean \pm SEM. Statistical significance was tested by the Student's *t*-test. Differences at the 95% level were considered significant.

Results

Animal characteristics

Data are summarized in Table 1. Five weeks after surgery the mean blood pressure was 178 mm Hg in 1K1C rats and 175 mm Hg in 2K1C rats; the development of blood pressure is shown in

Figure 1. In 1K1C hypertensive rats there was a significant decrease in water intake, hematocrit, plasma sodium concentration and plasma osmolality; plasma ANF and plasma norepinephrine concentrations were significantly increased. In contrast, 2K1C-hypertensive rats showed a significant increase in water intake, urine volume and hematocrit. The measurement of different components of the renin-angiotensin-aldosterone system revealed an expressed stimulation of this system with an about fivefold increase of plasma renin activity, plasma angiotensin II and plasma aldosterone concentration. Furthermore, plasma arginine vasopressin concentration was significantly increased.

ANF concentration in specific brain areas

The comparison of ANF concentrations in renovascular hypertensive rats with their normotensive control rats showed significant differences in nine brain areas of 1K1C hypertensive rats and in eight brain areas of 2K1C hypertensive rats (Table 2). In 1K1C hypertensive rats ANF concentration was significantly decreased only in the organum vasculosum laminae terminalis (OVLT) (28% of control). In all other brain areas with changes ANF concentration was significantly increased. Very pronounced elevations were found in the subfornical organ (211%), the supraoptic (244%) and paraventricular (172%) nuclei, the arcuate nucleus (245%), the median eminence (231%), the periventricular preoptic nucleus (169%) and the locus coeruleus (206%). The increase in the dorsal raphe nucleus (134%) was just moderate. In contrast, in 2K1C hypertensive rats ANF concentrations were increased only in the supraoptic nucleus (202%) and the medial amygdaloid nucleus (179%), whereas a significant decrease was found in the cerebral cortex (52%), the OVLT (36%), the periventricular preoptic nucleus (33%), the nucleus interstitialis striae terminalis (NIST) (59%), the median eminence (60%) and the medial forebrain bundle (69%).

Discussion

Present data confirm that there are differences in the pathogenesis of 1K1C and 2K1C hypertension. In 1K1C rats we have




Fig. 1. Development of blood pressure in 1K1C and 2K1C renovascular hypertensive rats and their respective control rats. Symbols are: (●) 1K1C on top, 1 kidney controls on bottom line; (○) 2K1C on top, 2 kidney controls on bottom dashed line.

Table 2. Concentrations of ANF in selected brain areas of 1K1C and 2K1C renal hypertensive rats and their control rats (pg ANF/mg protein; mean \pm SEM)

Brain area	1K1C control	1K1C	2K1C control	2K1C
Cerebral cortex	46.2 \pm 7.8	42.4 \pm 6.1	42.7 \pm 4.5	22.2 \pm 2.1 ^c
Org. vasc. lam. term.	83.0 \pm 7.0	23.0 \pm 1.7 ^c	107.1 \pm 3.7	38.7 \pm 6.5 ^c
Subfornical organ	73.6 \pm 6.8	155.6 \pm 8.7 ^b	76.6 \pm 15.8	86.0 \pm 19.8
Periventr. preopt. nucl.	337.5 \pm 16.0	570.8 \pm 75.8 ^a	400.6 \pm 47.3	134.2 \pm 18.3 ^c
Medial preopt. nucl.	237.8 \pm 9.4	235.0 \pm 20.0	219.6 \pm 35.3	173.1 \pm 12.8
NIST	140.6 \pm 20.9	214.8 \pm 21.8	123.7 \pm 13.3	73.3 \pm 7.2 ^b
Periventr. hypoth. nucl.	256.1 \pm 50.3	182.2 \pm 26.7	190.6 \pm 22.6	173.6 \pm 9.9
Supraoptic nucl.	44.2 \pm 8.1	107.8 \pm 9.0 ^b	47.9 \pm 5.9	96.8 \pm 10.4 ^b
Paraventricular nucl.	146.0 \pm 13.1	251.7 \pm 15.7 ^b	153.9 \pm 9.2	177.0 \pm 19.1
Arcuate nucl.	125.8 \pm 12.8	308.6 \pm 42.4 ^c	109.1 \pm 16.3	116.5 \pm 19.6
Median eminence	175.9 \pm 15.3	406.0 \pm 69.2 ^b	152.5 \pm 13.5	91.0 \pm 7.7 ^b
Perifornical nucl.	99.5 \pm 16.6	106.1 \pm 23.3	77.9 \pm 7.2	95.4 \pm 9.6
Median forebrain bundle	62.6 \pm 6.6	49.0 \pm 6.2	61.4 \pm 4.0	42.4 \pm 4.5 ^a
Medial amygd. nucl.	42.3 \pm 4.0	63.7 \pm 12.1	31.3 \pm 4.3	56.2 \pm 9.4 ^a
Dorsal raphe nucl.	90.6 \pm 12.2	121.2 \pm 8.7 ^a	104.3 \pm 14.9	115.0 \pm 24.7
Tegmentum pontis	118.6 \pm 21.6	70.3 \pm 11.6	95.4 \pm 10.0	83.9 \pm 6.3
Locus coeruleus	62.9 \pm 9.4	129.8 \pm 14.9 ^b	79.9 \pm 3.8	82.8 \pm 3.3
Nucl. tractus solitarii	74.3 \pm 5.6	65.0 \pm 8.3	54.2 \pm 4.1	51.6 \pm 3.8

^a $P < 0.05$, ^b $P < 0.01$, ^c $P < 0.001$, significant difference from the renovascular hypertensive rats to their respective control rats

found a normal plasma renin activity but a significantly increased plasma ANF level, which might be elevated in response to the augmented plasma volume in these animals. In contrast, the hypertension in our 2K1C model seems to be clearly renin dependent, since we have found an almost fivefold increase of plasma-renin-activity, plasma angiotensin II and plasma aldosterone concentration. The unchanged plasma ANF level and the significantly increased water intake tell against a relevant volume load in these animals.

The determination of ANF concentration in specific brain areas shows differential changes in 1K1C and 2K1C hypertensive rats. Except the unidirectional changes in the organum vasculosum laminae terminalis and in the supraoptic nucleus, ANF is changed in the opposite direction in all other brain areas, with an activation of the central ANF system in volume-

dependent 1K1C hypertension and an inhibition in renin-dependent 2K1C hypertension. In both models significant alterations were found in the circumventricular organs, in areas of the anteroventral third ventricle (AV3V) region and in the hypothalamohypophyseal system. In 1K1C hypertensive rats ANF was furthermore elevated in brain areas which are involved in the regulation of the sympathetic nervous system.

Of special interest are the alterations in the two investigated circumventricular organs, the subfornical organ (SFO) and the organum vasculosum laminae terminalis (OVLT). These brain areas lack a blood brain barrier and are exposed to changes in levels of circulating peptides such as ANF and may therefore represent the brain sites of action for circulating ANF. The SFO has long been implicated in the central regulation of electrolyte and fluid homeostasis and blood pressure [35–37]. In addition to

very high numbers of ANF receptors it contains also very high concentrations of angiotensin II (Ang II) receptors and has been shown to be the main brain target site for circulating Ang II [38]. Stimulation of Ang II receptors results in increased thirst, increased vasopressin secretion and increased blood pressure [39]. Since these actions are opposed by ANF [17–20,40] a local antagonism between these two peptides was postulated to exist in the SFO. In this regard, it is conceivable that the SFO might also be involved in the pathomechanism of renovascular hypertension. The increased binding of ANF in the SFO of 1K1C hypertensive rats could be a mechanism by which volume overload is signalled to intracerebral regulatory systems. In contrast, the unchanged ANF concentration in the SFO of 2K1C rats indicates that an imbalance between ANF and Ang II might exist there in these rats, which could be transferred to structures inside the blood brain barrier.

The OVLT contains not only ANF receptors, but also large numbers of ANF-immunoreactive nerve terminals and cell bodies. The coexistence of ANF receptors and ANF containing terminals outside the blood brain barrier suggests the possibility of an intimate connection between the peripheral and the central peptide system in the OVLT. Topographically and functionally, the OVLT belongs to the so-called anteroventral third ventricle (AV3V) region [41], which includes additionally the preoptic and hypothalamic periventricular nuclei and a part of the medial preoptic nucleus. This region incorporates the largest accumulation of ANF containing cells in the brain [21, 34]. It is sensitive to osmotic stimuli [41] and known to integrate, at a central level, mechanisms which modulate complex homeostatic systems, such as those associated with body fluid and electrolyte balance as well as cardiovascular regulation [41,42]. The importance of this brain region for the development of renovascular [15,16], Dahl [43] and deoxycorticosterone acetate salt [15] hypertension is well known. The changes of ANF concentration in the AV3V region of our models imply a special role for ANF mediated mechanisms at this level. In the OVLT, ANF is significantly decreased in both 1K1C and 2K1C hypertensive rats. Since a similar decrease was also found in spontaneously hypertensive rats [22], we suggest that ANF is decreased there as a consequence of the raised blood pressure. However, the mechanism remains to be elucidated.

Significant changes of ANF concentration have also been found in the hypothalamic supraoptic (SON) and paraventricular nuclei (PVN), where vasopressin is produced. On the basis of neuroanatomical and electrophysiological studies [19,44–48] it was suggested that vasopressin secretion is integrated by a regulatory unit which is composed of structures in the SFO, the AV3V region, the SON and PVN and uses Ang II and ANF as antagonistic neurotransmitters. The changes of ANF concentration in this regulatory unit indicate an altered influence on vasopressin secretion in both models of renovascular hypertension. But the significantly increased ANF level in 1K1C and 2K1C rats may be caused by different mechanisms. In volume loaded 1K1C rats ANF may be elevated there to inhibit vasopressin secretion. However, the unchanged plasma vasopressin concentration indicates that vasopressin secretion may still be too high for the state of volume dependent hypertension. In renin-dependent 2K1C rats ANF may be increased there to counteract the Ang II-stimulated vasopressin secretion. Interestingly, ANF in the PVN is only increased in 1K1C rats. This

may be explained by the different functional activity of the PVN and SON. While vasopressin and oxytocin secretion is dominant in the SON, the function of the PVN is more complex. Only the magnocellular subdivision of the PVN is engaged in vasopressin and oxytocin secretion, but the parvocellular subdivision contains several other neurotransmitters and neurohormones which are mainly involved in the regulation of the anterior pituitary gland [49]. Furthermore, cells in the PVN are integrated in a neural network with structures in the brain stem and the spinal cord and thereby involved in the regulation of blood pressure [50]. The increased ANF content in the PVN of 1K1C rats might therefore be related with these functions.

In this context it is of interest, that in the locus coeruleus ANF is increased only in 1K1C rats. The locus coeruleus is the quantitatively most important noradrenergic cell group of the brain and known to be involved in blood pressure regulation by modulation of the sympathetic nervous activity. It was shown that ANF can inhibit the increased sympathetic nervous activity in 1K1C hypertensive rats [51]. Therefore, we conclude that the increased ANF concentration in the locus coeruleus aims at an inhibition of the sympathetic nervous activity at this level and thereby counteracts the increased blood pressure. Interestingly, ANF is also increased in the locus coeruleus of DOCA-salt hypertensive rats; this form of hypertension is also volume dependent [38].

In conclusion, the central ANF system is differentially activated in 1K1C and 2K1C renovascular hypertension. In 1K1C hypertension, the activation of the central ANF system may be a compensatory mechanism to prevent further increments in blood pressure and plasma volume. In contrast, the depressed activity of the central ANF system in 2K1C hypertension may be a factor by which the elevation of blood pressure is promoted.

Acknowledgments

Preliminary results of this study have been presented as an oral presentation at the Fourth Scientific Meeting of The American Society of Hypertension, New York, 1989. We thank Mrs. Hilve Bonner, Miss Andrea Jahn and Mr. Jürgen Hupe for skilful technical assistance and Miss Christiane Winter for secretarial help.

Reprint requests to Udo Bahner, M.D., Medical Clinic of the University, Division of Nephrology, Josef-Schneider-Str. 2, D-8700 Würzburg, Federal Republic of Germany.

References

1. GOLDBLATT H, LYNCH J, HANZAL RF, SUMMERVILLE WW: Studies on experimental hypertension. I. The production of persistent elevation of systolic blood pressure by means of renal ischemia. *J Exp Med* 59:347–379, 1934
2. SEN S, SMEBY RR, BUMPUS FM, TURCOTTE JG: Role of renin-angiotensin system in chronic renal hypertensive rats. *Hypertension* 1:427–434, 1979
3. DE FORREST JM, KNAPPENBERGER RC, ANTONACCIO MJ, FERONE RA, CREEKMORE JS: Angiotensin II is a necessary component for the development of hypertension in the two-kidney, one-clip rat. *Am J Cardiol* 49:1515–1517, 1982
4. BRUNNER HR, KIRSHMAN JD, SEALEY JE, LARAGH JH: Hypertension of renal origin; evidence for two different mechanisms. *Science* 174:1344–1346, 1971
5. ALBERTINI R, BINIA A, OTSUKA Y, CARRETERO OA: Exchangeable sodium in angiotensinogenic and non-angiotensinogenic renovascular hypertension. *Hypertension* 1:624–630, 1979

6. MAACK T, CAMARGO MJF, KLEINERT HD, LARAGH JH, ATLAS SA: Atrial natriuretic factor: Structure and functional properties. *Kidney Int* 27:607-615, 1985
7. BALLERMAN BJ, BRENNER BM: Role of atrial peptides in body fluid homeostasis. *Circ Res* 58:619-630, 1986
8. NEEDLEMAN P, GREENWALD JE: Atriopeptin: A cardiac hormone intimately involved in fluid, electrolyte and blood pressure homeostasis. *N Engl J Med* 314:828-834, 1986
9. GARCIA R, CANTIN M, GUTKOWSKA J, THIBAUT G: Atrial natriuretic factor during development and reversal of one-kidney, one clip hypertension. *Hypertension* 9:144-149, 1987
10. GARCIA R, GUTKOWSKA J, GENEST J, CANTIN M, THIBAUT G: Reduction of blood pressure and increased diuresis and natriuresis during chronic infusion of atrial natriuretic factor (ANF Arg 101-Tyr 126) in conscious one-kidney, one-clip hypertensive rats. *Proc Soc Exp Biol Med* 179:539-545, 1985
11. GARCIA R, GUTKOWSKA J, CANTIN M, THIBAUT G: Renin dependency of the effect of chronically administered atrial natriuretic factor in two-kidney, one-clip rats. *Hypertension* 9:88-95, 1987
12. GARCIA R, THIBAUT G, GUTKOWSKA J, CANTIN M: Effect of chronic infusion of atrial natriuretic factor on plasma and urinary aldosterone, plasma renin activity, blood pressure and sodium excretion in 2K1C hypertensive rats. *Clin Exp Hypertens* A8:1127-1147, 1986
13. REID JL, DARGIE HJ, FRANKLIN SS, FRASER B: Plasma noradrenaline and renovascular hypertension in the rat. *Clin Sci Mol Med* 51:439s-442s, 1976
14. DE QUATTRO V, EIDE I, MYERS MR, EIDE K, KOLLACH R, WHIGHAM H: Enhanced hypothalamic noradrenaline biosynthesis in Goldblatt I renovascular hypertension. *Clin Sci Mol Med* 55:109-112, 1978
15. BUGGY J, FINK GD, JOHNSON AK, BRODY MJ: The effect of anteroventral third ventricle tissue lesions on the development of renal hypertension in rats. *Circ Res* 40:1110-1117, 1977
16. HAYWOOD JR, FINK GD, BUGGY J, BONTELE S, JOHNSON AK, BRODY MJ: Prevention of two-kidney one-clip renal hypertension in the rat by ablation of anteroventral third ventricle (AV3V) tissue. *Am J Physiol* 245:H683-H689, 1983
17. FITTS DA, THUNHURST RL, SIMPSON JB: Diuresis and reduction of salt appetite by lateral ventricular infusions of atriopeptin II. *Brain Res* 348:118-124, 1985
18. ANTUNES-RODRIGUES J, MCCANN SM, SAMSON WK: Central administration of atrial natriuretic factor inhibits saline preference in the rat. *Endocrinology* 118:1726-1728, 1986
19. IITAKE K, SHARE L, CROFTON JT, BROOKS DP, OUCHI Y, BLAINE EH: Central atrial natriuretic factor reduces vasopressin secretion in the rat. *Endocrinology* 119:438-440, 1986
20. ISRAEL A, BARBELLA Y: Diuretic and natriuretic action of rat atrial natriuretic peptide (6-33) administered intracerebroventricularly in rats. *Brain Res Bull* 18:141-144, 1986
21. STANDAERT DG, NEEDLEMAN P, SAPER CB: Organization of atriopeptin-like immunoreactive neurons in the central nervous system of the rat. *J Comp Neurol* 253:315-341, 1986
22. BAHNER U, GEIGER H, PALKOVITS M, HEIDLAND A: Atrial natriuretic factor in specific brain areas of spontaneously hypertensive rats. *Hypertension* 12:519-524, 1988
23. GEIGER H, BAHNER U, PALKOVITS M, HUPE J, HEIDLAND A: Atrial natriuretic factor content of brain nuclei in deoxycorticosterone acetate-salt hypertension in the rat. *Clin Sci* 77:529-534, 1989
24. BAHNER U, GEIGER H, PALKOVITS M, GANTEN D, MICHEL J, HEIDLAND A: Atrial natriuretic peptides in brain nuclei of rats with inherited diabetes insipidus (Brattleboro rats). *Neuroendocrinology* 51:721-729, 1990
25. GANTEN D, SCHELLING P, HOFFMANN WE, PHILLIPS MI, GANTEN U: The measurement of extrarenal isorenins; in *Radioimmunoassay-renin-angiotensin*, edited by KRAUSE DK, HUMMERICH W, POULSON K, Stuttgart, Thieme, 1978, pp. 144-155
26. MANN JFE, JOHNSON AK, GANTEN D: Plasma angiotensin II: Dipsogenic levels and angiotensin-generating capacity of renin. *Am J Physiol* 238:R372-R377, 1980
27. RASCHER W, WEIDMANN E, GROSS F: Vasopressin in the plasma of stroke-prone spontaneously hypertensive rats. *Clin Sci* 61:295-298, 1981
28. KRSTULOVIC AM: *Quantitative analysis of Catecholamines and Related Compounds*. New York, Chichester, Brisbane, Toronto, John Wiley and Sons, 1986
29. LAROSE P, MELOCHE S, DU SONICH P, DELEAN A, ONG H: Radioimmunoassay of atrial natriuretic factor: Human plasma levels. *Biochem Biophys Res Commun* 130:553-558, 1985
30. PALKOVITS M: Isolated removal of hypothalamic or other brain nuclei of the rat. *Brain Res* 59:449-450, 1973
31. LOWRY OH, ROSEBROUGH NY, FARR AL, RANDAL RJ: Protein measurement with the Folin phenol reagent. *J Biol Chem* 193:265-275, 1951
32. SCHWARTZ D, GELLER DM, MANNING PT, SIEGEL NR, FOK KR, SMITH CE, NEEDLEMAN P: Ser-Leu-Arg-Arg-atriopeptin III: the major circulating form of atrial peptide. *Science* 229:397-400, 1985
33. SHONO S, NAKAO K, MORII N, YAMADA T, ITOH H, SAKAMOTO M, SUGAWARA A, SAITO Y, KATSUURA G, IMURA H: Nature of atrial natriuretic polypeptide in rat brain. *Biochem Biophys Res Commun* 135:728-734, 1986
34. ZAMIR N, SKOFITSCH G, ESKAY RL, JACOBOWITZ DM: Distribution of immunoreactive atrial natriuretic peptides in the central nervous system of the rat. *Brain Res* 365:105-111, 1986
35. PALKOVITS M: The role of the subfornical organ in the salt water balance. *Naturwissenschaften* 13:336-337, 1966
36. SIMPSON JB, ROUTTENBERG A: Subfornical organ: Site of drinking elicitation by angiotensin. *Science* 81:1172-1175, 1973
37. MANGIAPANE ML, SIMPSON JB: Subfornical organ: Forebrain site of pressor and dipsogenic action of angiotensin II. *Am J Physiol* 239:R382-R389, 1980
38. MENDELSON FAO, QUIRION R, SAAVEDRA JM, AGUILERA G, CATT K: Autoradiographic localization of angiotensin II receptors in rat brain. *Proc Natl Acad Sci USA* 81:1575-1579, 1984
39. PHILLIPS MI: Functions of angiotensin in the central nervous system. *Annu Rev Physiol* 49:413-435, 1987
40. MASOTTO C, NEGRO-VILAR A: Inhibition of spontaneous or angiotensin II-stimulated water intake by atrial natriuretic factor. *Brain Res Bull* 15:523-526, 1985
41. BRODY MJ, JOHNSON AK: Role of the anteroventral third ventricle region in fluid and electrolyte balance, arterial pressure regulation and hypertension, in *Frontiers in Neuroendocrinology* (Vol 6) edited by MARTINI L, GANONG WF, New York, Raven Press, 1980, pp. 249-292
42. HARTLE DK, BRODY MJ: The angiotensin II pressor system of the rat forebrain. *Circ Res* 54:355-366, 1984
43. GOTO A, GANGULI M, TOBIAN L, JOHNSON MA, TWAI J: Effect of an anteroventral third ventricle lesion on NaCl hypertension in Dahl salt-sensitive rats. *Am J Physiol* 243:H614-H618, 1982
44. MISELIS RR: The efferent projections of the subfornical organ of the rat: A circumventricular organ within a neural network subserving water balance. *Brain Res* 230:1-23, 1981
45. SAWCHENKO PE, SWANSON LE: The organization of forebrain afferents to the paraventricular and supraoptic nuclei of the rat. *J Comp Neurol* 218:121-144, 1983
46. FERGUSON AV, DAY TA, RENAUD LP: Subfornical organ efferents influence the excitability of neurohypophyseal and tuberoinfundibular paraventricular nucleus neurons in the rat. *Neuroendocrinology* 39:423-428, 1984
47. SAMSON WK: Atrial natriuretic factor inhibits dehydration and hemorrhage-induced vasopressin release. *Neuroendocrinology* 40:277-279, 1985
48. STANDAERT DG, CECETTO DF, NEEDLEMAN P, SAPER CP: Inhibition of the firing of vasopressin neurons by atriopeptin. *Nature* 329:151-153, 1987
49. PALKOVITS M, ESKAY RL, ANTONI FA: Atrial natriuretic peptide in the median eminence is of paraventricular nucleus origin. *Neuroendocrinology* 46:542-544, 1987
50. ZERIHUN L, HARRIS MC: An electrophysiological analysis of caudally-projecting neurones from the hypothalamic paraventricular nucleus in the rat. *Brain Res* 261:13-20, 1983
51. DEBINSKI W, KUCHEL O, GARCIA R, BUN NT, RACZ K, CANTIN M, GENEST J: Atrial natriuretic factor inhibits the sympathetic nervous activity in one-kidney, one-clip hypertension in the rat. *Proc Soc Exp Biol Med* 181:173-177, 1986